

Kirkland Lake

53 Government Rd. W./
53, ch. Government O.
Kirkland Lake ON P2N 2E5

Tel/Tél : 705 567 2222
Fax/Télex : 705 567 9407

North Bay

1164 Devonshire Ave./
1164, av. Devonshire
North Bay ON P1B 6X7

Tel/Tél : 705 476 2222
Fax/Télex : 705 474 0080

Parry Sound

6 Albert St./
6, rue Albert
Parry Sound ON P2A 3A4

Tel/Tél : 1 800 440 6762
Fax/Télex : 1-855-773-4056

Sault Ste. Marie

390 Bay St., Main Floor/
390, rue Bay, 1e étage
Sault Ste. Marie ON P6A 1X2

Tel/Tél : 705 949 1650
Fax/Télex : 705 949 1663

Sudbury

40 Elm St., Suite 41-C/
40, rue Elm, bureau 41-C
Sudbury ON P3C 1S8

Tel/Tél : 705 522 3461
Fax/Télex : 705 522 3855

Timmins

330 Second Ave., Suite 101/
330 av. Second, bureau 101
Timmins ON P4N 8A4

Tel/Tél : 705 267 7766
Fax/Télex : 705 267 7795

TTY / ATS 711 (ask operator for 1-888-533-2222 / veuillez demander le téléphoniste pour le 1-888-533-222)
Toll Free / Sans frais: 1-800-461-2919 or / ou 310-2222 no area code required / indicatif régional non requis.

Negative Pressure Wound Therapy (NPWT) Referral Form

PATIENT INFORMATION			
Name:		Health Card No and Version Code:	
Address:		Postal Code:	
Date of Birth (dd-mmm-yyyy):		Phone:	
Gender: ☐ Male ☐ Female ☐ Non-binary ☐ Unknown		Pronouns:	
Diagnosis:		Diabetic: ☐ Yes ☐ No	
Allergies: ☐ Yes ☐ No ☐ Unknown Specify:		Latex Allergy: ☐ Yes ☐ No ☐ Unknown	
WOUND TYPE			
The following conditions can be considered for the application of NPWT. Please indicate reason for referral.			
Acute Wound	☐ Surgical (dehiscid) ☐ Traumatic ☐ Abdominal ☐ Pilonidal cyst ☐ Partial thickness burn		
Chronic Open Wound	☐ Diabetic ulcer (offloaded) ☐ Venous leg ulcer ☐ Stage 3 or 4 pressure injury (offloaded)		
Adjunct to Surgery	☐ Preparation of wound bed post-operatively ☐ Incisional support ☐ Securing skin graft post-operatively		
Oncology Related	☐ Wound complicated by radiation ☐ Support wound healing prior to start of chemotherapy		
WOUND DESCRIPTION			
Location:		Length: cm x Width: cm x Depth: cm	
☐ Undermining Details if applicable:		☐ Tunneling Details if applicable:	
Note: NPWT will continue to be assessed in the community, and settings may be reviewed based on exudate and patient tolerance. Continuation of NPWT is dependent on wound healing goals being met. Maximum treatment time for NPWT is 8 weeks.			
NPWT TREATMENT ORDERS			
☐ ActiVAC (indicate pressure settings and dressing details below) Pressure (mmHg): ☐ Continuous OR ☐ Intermittent Dressing (select all that apply) Granufilm Black: ☐ Small (10 cm x 7.5 cm x 3.2 cm) ☐ Medium (18 cm x 12.5 cm x 3.2 cm) ☐ Large (26 cm x 15 cm x 3.2 cm) ☐ X-Large (60 cm x 30 cm x 3.2 cm) Granufilm Silver: ☐ Small (10 cm x 7.5 cm x 3.2 cm) ☐ Medium (18 cm x 12.5 cm x 3.2 cm) ☐ Large (26 cm x 15 cm x 3.2 cm) White Foam: ☐ Small (10 cm x 7.5 cm x 1 cm) ☐ Large (10 cm x 15 cm x 1 cm) Peel and Place: * Peel and Place cannot be used with tunneling or undermining > 2cm		☐ PICO (single use, disposable) Pressure: ☐ 80 mmHg (non-adjustable) Dressing Size: ☐ 10 cm x 20 cm ☐ 10 cm x 30 cm ☐ 15 cm x 15 cm ☐ 15 cm x 20 cm ☐ 15 cm x 30 cm ☐ 20 cm x 20 cm	
		☐ SNAP (single use, disposable) Pressure: ☐ 125 mmHg (non-adjustable) Dressing Size: ☐ 10 cm x 10 cm ☐ 15 cm x 15 cm	

☐ Small (wound size less than 6.1 x 8.6 cm, MAX 2 cm deep) ☐ Medium (wound size less than 11.1 x 16.6 cm, MAX 4 cm deep) ☐ Large (wound size less than 13.6 x 24.2 cm, MAX 6 cm deep)	SNAP Therapy Strap Size: ☐ Small ☐ Medium ☐ Large
--	---

Patient Name:	HCN:
----------------------	-------------

CONVENTIONAL DRESSING ORDERS

Patients will be started on conventional dressings until NPWT can be initiated. Conventional orders also required in the case of service interruption.

PRECAUTIONS AND CONTRAINDICATIONS
--

The precautions and contraindications listed below have been reviewed, and its determined that NPWT is appropriate to be used for patient
☐ **Yes** ☐ **No** (conventional dressings will be utilized until addressed)

The following conditions are considered precautions in the use of NPWT: • Immunodeficiency (e.g. Leukemia, HIV); • Hematologic disorders; • Systemic or local signs of infection; • Uncontrolled diabetes; • Systemic steroids; • Receiving anticoagulant therapy; • The location of the wound will interfere with the therapy; • Nutritional impairment; • History of non-compliance; • Home environment not conducive to NPWT (i.e. cleanliness, animals, etc.); or • Patient unable to adhere to minimum of 22 hours of therapy/day.	The following risk factors contraindicate the use of NPWT; • Inadequate wound visualization; • Untreated infection in the wound site; • Fistulas to body cavities or organs; • Presence of undebrided necrotic tissue with eschar; • Untreated Osteomyelitis; • Malignancy or cancer in the wound margins; • Unresolved bleeding following debridement; or • Exposed vasculature, nerves or organ
---	--

PRESCRIBER INFORMATION

Name:	Phone:	Fax:
Signature:	After Hours Number:	
CPSO/CNO#:	Date:	